Lee County Telephone (229) 759-6025
P. O. Box 589 Fax (229) 759-8302
Leesburg, Georgia 31763 uge4177@uga.edu

Lee County Ag Newsletter

September 2022, Volume 22, Number 11

Water Requirement vs Water Supplied (a different perspective)

R. Scott Tubbs & Wesley M. Porter, UGA

tZ v }v•] Œ]vP u}všZoÇ Œ]v(oo À Œ P •U o}}l• v]À]vPX &}Œ Æ u current 2022 growing seas at Midville, GA according to the University of Georgia Weather Network (georgiaweather.net). I June. Although scenario C shows

Monthly Rainfall (season total = 16.60 inches)

	May 1-31	June 1-30	July 1-31	Aug 1-26
	3.27	2.74	5.60	5.08

	Apr 28-May 28	May 29-June 27	June 28-July 28	July 29-Aug 26

Figure 1. Water requirements of late April planted peanuts with 1.25 average and actual rainfall.

If we shift this same concept to a peanut crop that was planted late in the planting window (i.e. the first week of June), th
Œ } ‰ [• Á š Œ v • } v } š u š Z š Z Z] • š } Œ] o À Œ P Œ] v (o o • o } • o Ç X
rainfall is typically becoming more scarce.

& } Œ š µ v š o Ç U š Z µ Œ Œ v š Œ] v (o o Œ] À µ ‰ µ v š] o š Z] • ‰ }] v š] v a l š Z a î ì î

Figure 2. Water requirements of early June planted peanuts with 1.25 y average and actual rainfall.

It should also be noted that the checkbook curves shown above are a general guide to demonstrate how water demand increases, peaks, then decreases over the season. However, many other factors are involved in more accurately determini the localized needs of a given field, including soil type/texture, organic matter content, porosity/drainage, and other soil health Z Œ š Œ] • š] • X À v • Z µ o] v P š Z v] ‹ µ • Z À v À o } ‰] v Œ on v š requirements. There are smartphone apps, web-based schedulers, soil moisture sensors, etc. that are more precise in targetin š µ o] Œ Œ] P š] } v Œ ‹ µ] Œ u v š • X d Z • v À Œ Ç µ • (µ o] v u š] v P š Z Œ maximum yield potential.

End of Season Irrigation for Peanuts David Hall, Jason Mallard, and Wesley Porter, UGA

The only thing that is consistent from year to year is that each season is different and variable. While last year had high of rainfall, this year has been very hot and dry from May through the end of June. Since the end of June, we have been getting sporadic rainfall across parts of the state. Thus, you need to monitor what your current soil moisture condition is and make appropriate decisions moving forward. Additionally, keep an eye on the long term forecast and the tropics. Up to this point, we have been getting the sporadic rainfall, but we have been lucky that there have not been any significant tropical events. Tha

would want to do is scheduleirrigation without boots on the ground or moisture sensors relaying real time data, therefore risking increased disease outbreaks or soil drying out. During peak water demand and dry weather, it is fairly simple to sche irrigation events. This time of the season water demand begins to fall off quickly and most have been receiving ample rainfall. Do not let your guard down if we enter a dry period with dry hot west winds. (We prefer those conditions after digging!) Sand soils can dry out fast and we are looking at what appears to be a great crop. Remember, heavy downpours that exceed the so water holding capacities basically become run off. We have received much of that this year.

The month of September is when the majority of our peanuts are dug and most of them are now well past the peak water demand and need less than an inch of water per week. Now is a good time to start thinking about irrigation termination for

Points to Consider for Late-Season Disease Control in Peanuts
Bob Kemerait, UGA

Late-season disease recommendations for a peanut crop are often confusing. Reasons include A)] P P] v P l Z Œ À • š š • Œ v } š ^ • š] v • š } v _ U • š Z] v] v } (} v] • fungicides, C) disease can be cryptic, as in the case of underground white mold, D) there are a number of fungici options that can be deployed, and E) approaching rains can make it necessary to quickly change management pla t Z] o š Z Œ u Ç v } š •] v P o ^ • š _ Œ } u u v š] } v U • } u • } o µ š] } v • Œ have three things in common: a) timeliness, b) use of the right products (emphasis that there is often more than one ^ Œ] P Z š _ ‰ Œ } µ š • U v • µ • } (š Z Œ] P Z š ‰ Œ } management decisions are made, and c) more] u ‰ } Œ š v š v } Á š Z v š Z Ç À Œ Z À v • u u Z } (} Œ P] [• ‰ Ç • X K µ Œ Z] • š } Œ] ^ • ‰ Œ Ç À Œ Ç í ð Ç • (} Œ š } š o } (ó • ‰ Œ Ç • _ u

Growers often request advice on adjusting digging dates based upon disease in the field. Generally, it is best to until harvest maturity is reached in order to assure maximum grade, rather than digging the peanuts early. For example, though tomato spotted wilt may be severe in a field in 2022, I generally recommend waiting until harves maturity to dig the peanut U µ v o • • } š Z Œ] • • • U o] l -Á Z] š v P u } o } v U š } affected by the tomato spotted wilt. However, where defoliation from leaf spot is severe, as it is in some field now then it may be critical to digging earlier than planned in order to protect yield. Georgia-06G can withstand defoliation of 50-60% IF digging is NOT delayed past maturity. Any unexpected delays in digging will result in yiel losses. Where white mold is severe, for example greater than 50% incidence, the grower should consider digging early. Significant defoliation from leaf spot diseases and severe outbreaks of white mold can increase digging los by weakening pegs.

NOTE: A critical consideration for choice of fungicides late in the season is that pre-harvest intervals (PHI) vary among fungicides. For example, Alto has a 30-day PHI, and Elatus and Convoy have 40-day PHI, compared to 14-day PHI for other fungicides such as Provost Silver and Fontelis. Growers must always check the label to make sure on all of these.

Below are some typical situations that peanut growers may find themselves in and suggestions for control:

Grower is 4 or more weeks away from harvest and currently has excellent disease control.

- Suggestion: I recommend the grower apply at least one more fungicide for leaf spot control with an inexpensive white mold material mixed with it, for good measure. It is generally helpful to use a mix of a protectant leaf spot fungicide (like chlorothalonil) mixed with a leaf spot fungicide with some curative activity (e.g., Alto, Domark, Topsin) for best protection.
- Suggestion: Given the low cost of tebuconazole, the grower may consider applying a mix of tebuconazole + chlorothalonil for added insurance of white mold and leaf spot control. NOTE 1: If white mold is not an issue, then the grower may stick with a leaf spot spray only. Note 2: If grower has planted Georgia-06G or Georgia-12Y and the plants are leaf spot-free at 4 weeks prior to the anticipated digging date, an additional fungicide application for leaf spot may not be needed if grower is willing to watch/scout the field for oth

expected U.S. cotton production. In 2022, the U.S. planted 12.3 million acres of upland cotton, the highest in 3 years, which was mainly due to historically high cotton prices during the decision-making and planting window.

However, in 2022, the overall U.S. abandonment rate for upland cotton is estimated at 43.4%, which is the highest on record since 1953. The abandonment rate, which measures the percentage of unharvested acres compared to total planted acres, provides an estimate of the number of failed acres versus the number of acres that will be harvested. Severe drought conditions hit the largest cotton production regions in the Southwest (Texas, Oklahoma, and Kansas) and the West (California, Arizona, and New Mexico). The abandonment rate for Texas (Figure 1A) reached 69%. Texas planted 7.1 million acres of cotton in 2022 by far the largest of any state, representing 57.6% of total U.S. planted acres (Figure 1B). By contrast, drought impacts were less severe in the Delta (Missouri, Arkansas, Louisiana, Mississippi, and Tennessee) and Southeast (Alabama, Georgia, Florida, South Carolina, North Carolina, Virginia).

As a result of the drought conditions this year, upland cotton harvested acreage in the U.S. is projected at 7.0 million acres, which is the lowest amount of harvested acreage in over 150 years. The projected high abandonment rate in the U.S. reduced expected cotton production to 12.2 million bales, compared to the 10-year average of 16 million bales, according to

keep in mind that the cotton plant moisture uptake is low, but there still is a need for some moisture in the soil to aid in finishing out those last harvestable bolls. At this point if rainfall keeps up, there will be very little if no need for additional irrigation.



Figure 1. Seasonal cotton water usage with peak, decline and irrigation termination highlighted.

As water use declines, it is necessary to closely monitor your fields for cotton bolls that are beginning to R S H Q 8 * $ ¶ V R I I L F L D O L U U L J D W L R Q W H U P L Q D W L R Q U H F R P P H Q D W L R Q S R O L F \ across a field. 10% open is not a high number of bolls on a plant that are open to start considering irrigation termination. A plant with 12 harvestable bolls means you need to be looking at the available soil moisture if 1.2 bolls are open. September 2019 was abnormally hot and dry. Bolls were beginning to open but our available water in the soil was depleted very rapidly due to the environmental conditions. Due to dry, hot temps and long-term dry weather forecasts, one last irrigation event(s) was(were) needed to help finish the crop. The low humidity and sunny conditions helped to prevent open bolls from deteriorating. Currently, we are seeing high levels of moisture and boll rot similar to what we witnessed during 2020 because heavy dews, high humidity and plentiful rainfall are excellent conditions for it. Do not irrigate itself or worsen boll rot issues if sufficient soil moisture is present. In times when an application of irrigation is needed, soak the soil sufficiently so that you can limit your irrigation events to as few as possible. Frequent small irrigation events will only exacerbate or flare boll rot. Paying attention to the long term and short-

This boll value tree really emphasizes the importance of first setting those high dollar bolls early in the season with proper management. Nearing the end of the growing season, watering and irrigating your crop with a higher than 10% open boll in an effort to make young upper position low value bolls open in the top will lead to losing or damaging your most valuable bolls and reducing yield and profitability. Not to mention, those young bolls probably will not mature enough to be harvested anyway. If you have questions about finalizing irrigation make sure you reach out to your local UGA County Extension Agent.

Weather and Climate Outlook for September 2022 and Beyond (Pam Knox): Most of the Southeast experienced conditions in August that were wetter and warmer than usual, although as usual there were variations across the region. Some areas are still feeling the impacts of too much rain, while others have experienced dry conditions that have reduced yields in their crops. The humid conditions and lack of sunshine in the areas that have experienced a lot of rain are feeling the impacts in the form of abundant fungal diseases and a difficult time for farmers to get into their fields.

September so far looks like it will be a continuation of warm and wet conditions, although there will be some more seasonal periods and some areas. There will be some periods of dry weather scattered amid rainy days, so you will need to watch the weather forecasts carefully to find those windows of opportunity to work in the fields.

Later in fall, I expect to see drier conditions appear. This is not unusual in October, since this is the driest month of the year for many parts of the region, but it could be drier than usual as La Niña remains strong and continues to affect our weather over the fall and winter. Generally, La Niña has the strongest signal in southern Georgia and Alabama and into Florida, with more northern areas less predictable b

visual observations only. I cannot explain why

3. Agronomic crops such as cotton and soybean.
4. Fall vegetables which include a similar crop mix as spring vegetables.

Management of SLWF in these individual cropping systems affects infestations in subsequent crops as SLWF move from one cropping system to the next. Failing to properly manage SLWF in any one of these systems can have negative consequences for subsequent cropping systems. It is important that we manage SLWF in all cropping systems to manage the overall SLWF population. The SLWF population is somewhat like a snowball and just grows larger and larger during the summer. Have you ever considered how many SLWF are produced per acre if we defoliate 1, 2, or 3 weeks late? Timely defoliation will lower the risk of having fiber quality problems related to whitefli. Ultimately how we manage SLWF now will have impacts on us in 2023. By in large, Georgia growers have done a good job managing SLWF in cotton. Keeping in mind that SLWF requires management until harvest, one of the best practices is to



2. Make sure you are using the correct products and rates to accomplish your intended goals. There are 3 main goals of defoliation – leaf removal (juvenile and mature leaves), regrowth prevention, and boll opening. Determining which goals you are trying to accomplish, as well as the environmental conditions surrounding defoliation, will assist in the decision on products and rates. For regrowth control and juvenile leaf removal, thidiazuron containing products will be needed in the tank mix. Thidiazuron alone is available under many trade names (Dropp, Freefall, Klean-pik, etc.), and there is also a premix of thidiazuron-diuron (Ginstar, Cutout, Adios, etc.). Mature leaf removal can be accomplished with a number of products. The main one utilized in Georgia is Folex (tribufos), but there are a number of PPO inhibiting herbicides that can be utilized as well (Aim, ET, Spartan, Reviton, Display, Blizzard, Resource). Lastly, your boll opening products contain ethephon. Whether it is ethephon alone (Boll Buster, SuperBoll, Setup, etc.), ethephon + urea sulfate (Cotton Quik, First Pick), or ethephon + cyclanalide (Finish), with the goal of open bolls.

The most common tank-mixture utilized by Georgia growers includes thidiazuron (Dropp), tribufos (Folex), and ethephon (Prep). Below is a table to assist in rate selection for each product from early to late season.

For ethephon and tribufos, as the season progresses and it gets cooler, rates should increase to g the desired effects. Whereas once we get to the end of the season, thidiazuron is removed from the tank-mix because as it gets cooler outside there is a lower risk of regrowth. Keep in mind that you can substitute tribufos for any of the aforementioned PPO inhibiting herbicides, and I would follow the same trends with rate selection (lower rates when L W s r m e outside, increase as it gets cooler). Recommended rates on each product discussed here throughout the season can be found beginnin on page 148 of the 2022 UGA Cotton Production Guide, which is at the following link: http://www.ugacotton.com/productionguide/. Also, I will be in constant contact with our UGA County Extension agents throughout defoliation and harvest season, so if you have questions abou products or rates feel free to reach out to your local UGA County Extension Agent.

3. Zero Tolerance for regrowth in 2022!!! Dr. Phillip Roberts has already discussed this in his newsletter entry, but keep in mind that one of the best things you could do to manage whiteflies in 2022 and assist in management for 2023 is defoliate your cotton crop in a timely manner. Below is a picture I took in some plots in Tifton. It is clear that the bottom crop is ready to defoliate, but with recent rains it has started regrowing and blooming. While many would try to ^ ‰ µ š š } ‰, we decided to defoliate it. Investing more money to try to make a top crop } • v [š • u Á } Œ š Z] š (} Œ t first, we have enough money tied up in this crop as it is, and waiting on that top to finish will cause the bottom crop to deteriorate. Œ š X > š [• P š Á Z š [• š Z Œ v o o] š Ç Œ X ^ } v U] (Á o š š Z š Œ P Œ } A like cotton candy to whiteflies (circled in red below). They will keep feeding and reproducing there, and will contribute to the population going into the winter time and consequently into v Æ š Ç Œ X > š [• (} o] š ‰] v š] u o Ç

4. One shot vs. two shot what are the benefits? I get this question a good bit during defoliation time. In Georgia, we primarily use 1 shot for defoliation and it works very well. Rarely will I ever recommend a second shot unless something went wrong on the first. So keep in mind the amount of money already invested in this crop a second shot at defoliation means more chemical costs, as Á o o •] • o (µ o l ‰ ‰ o] š] } v } • š • X > š [• P š] š } get the crop out in a timely manner.

5. D } [Á š Œ] • u } [š š Œ X Z] P Z Œ • ‰ Œ Ç Œ } µ š ‰ µ š ~ P o o } v •

would like to submit a comment to the EPA stating how important these chemistries are to you andyour

operation, please contact your local UGA County Extension Agent. They all know how to do this, and I promise that the EPA reads these comments and values input from producers.

7. Timely defoliation and harvest are key to producing a high quality crop. Only defoliate what you can come back and harvest in about 10 to 14 days. This helps preserve crop quality and can reduce the likelihood of an extraneous matter call. And speaking of extraneous matter, handle your modules carefully and be sure to get grocery bags/balloons out of your field prior to harvest to reduce the likelihood of plastic

}vš u]v š]}vX /š[• îy µ š]}v (Œ}u šZ o ••]vP }(()] v

+ H U H ¶ V W K H Q H Z O L V W

Go to the UGA Variety testing website for yield and more info. https://swvt.uga.edu/



Ultra Late Soybeans Growing

+HUH¶V VRPH VR\EHDQV SODQWHG DIWHU First Field KDUY
was planted in 7.5 inch rows with a grain drill. Then we have a field planted with a 30 inch planter, and the last photo is one where they used the 30 inch planter but burned the corn stalks. 50 pounds of N was applied to get these all growing tall, fast.







Georgia Grain News 8-25-22

Corn Yield Test Results

+HUH¶V VRPH FRUQ K\EULG SORW \LHOG WHVW UHVXO
getting these done. In choosing hybrids to plant it is good to look at as much data as possible including state yield tests and any local data.

Grady/Decatur Corn Hy brid Test

Contact Cale Cloud in Grady county or Clark Higgins of Decatur county for more info.

Appling County Corn Hybrid Test



:LWK WKH H[FHSWLRQ RI VRPH VFDE SUHVVXUH WKURXJKRXW - conditions all season to this point and the appearance of the crop shows it. Growers have done a fine job of protecting the nuts from scab throughout the rainy periods. Industry forecasts currently have Georgia estimated at about 130 million lbs. But, as we have learned in years past, just EHFDXVH ZH FDQ VHH WKH ILQLVK OLQH GRHVQ¶W PHDQ ZH¶Y

We have been in a similar position as recently as last year when we had low solar radiation season-long. But, the most serious blow to the 2021 crop came from cloudy weather during the kernel filling stage. The sunniest part of the year in our part of the world usually runs from September 11-December 10. If you recall, crop development in 2021 was somewhat late. For these purposes, solar

Doug Collins
Lee County Extension Coordinator
P. O. Box 589
Leesburg, GA 31763
229 - 759 - 6025
229 - 759 - 3302 - FAX
229 - 344 - 8031 - cell
collinsd@uga.edu